


α -Carotene and β -carotene are a type of carotenoid and widely known as precursors to vitamin A in human body. They also have antioxidant activity.


The two carotenes are very similar in chemical structure. Since the difference is only in the position of a carbon-carbon double bond, it is difficult to separate them with widely-used monomeric ODS columns. To solve this problem, Inertsil ODS-P, which is a polymeric ODS column and superior in steric recognition, was used as a separation column in this note.

(K. Suzuki)



Chromatograms obtained from standard solution

When a commercially available monomeric column was used, α -carotene and β -carotene were not separated within 20 minutes. However, Inertsil ODS-P provided good separation of the two carotenes.



Conditions

Column : Inertsil ODS-P
(5 μ m, 250 x 4.6 mm I.D.)
Cat.No. 5020-02002


Eluent : A) CH₃OH
B) C₂H₅OH
A/B = 90/10, v/v
(gradient mixer)

Flow rate : 1.0 mL/min

Col. Temp. : 40 °C

Detection : PDA 455 nm

Inj. Vol. : 20 μ L



Analyses of food samples

Sample pretreatment 1

Vegetables

- 0.5 g
- 3 % Pyrogallol ethanol solution 10 mL
- 60 % KOH aqueous solution 1 mL

Saponification

- Heat at 70 °C for 30 min
- Place in an ice-water bath to cool down
- 1 % NaCl aqueous solution 22.5 mL
- Hexane-ethyl acetate = 9/1 (v/v) 15 mL
- Shaking for 5 min

Liquid-liquid extraction

- Organic layer
- Aqueous layer
- Repeat procedure *A twice

Concentration

- Evaporate under reduced pressure
- Dissolve in 2 mL of ethanol

HPLC

Sample pretreatment 2

Carrot juice

- 2 g
- 3 % Pyrogallol ethanol solution 20 mL
- Sodium sulfate anhydrous 10 g
- Shaking for 5 min

Centrifuge

- Supernatant
- Residue
- Repeat procedure *B twice

- 1 % NaCl aqueous solution 23 mL
- 2-Propanol 6 mL
- Hexane-ethyl acetate = 9/1 (v/v) 15 mL
- Shaking for 5 min



Liquid-liquid extraction

- Organic layer
- Aqueous layer
- Repeat procedure *A twice

Concentration

- Evaporate under reduced pressure
- Dissolve in 10 mL of ethanol

HPLC



GL Sciences disclaims any and all responsibility for any injury or damage which may be caused by this data directly or indirectly. We reserve the right to amend this information or data at any time and without any prior announcement.

GL Sciences Inc. Japan

22-1 Nishishinjuku 6-chome
Shinjuku-ku, Tokyo
163-1130, Japan

Phone: +81-3-5323-6620
Fax: +81-3-5323-6621
Email: world@glsc.co.jp
Web: www.glsciences.com

GL Sciences Inc. USA

4733 Torrance Blvd. Suite 255
Torrance, CA 90503
USA

Phone: +1-310-265-4424
Fax: +1-310-265-4425
Email: info@glsciencesinc.com
Web: www.glsciencesinc.com

GL Sciences B.V.

Dillenburgstraat 7C
5652AM, Eindhoven
The Netherlands

Phone: +31-40-254-9531
Email: info@glsciences.eu
Web: www.glsciences.eu

GL Sciences (Shanghai) Limited

Tower B, Room 2003
Far East International Plaza
No.317 Xianxia Road, Changning District
Shanghai, China 200051

Phone: +86-21-62782272
Email: contact@glsciences.com.cn
Web: www.glsciences.com.cn



International Distributors

Visit our Website at www.glsciences.com/distributors